



GOAT HEALTH TREATMENT CHECKLIST



Condition	Signs	First Aid	Medication	Follow-Up
Diarrhea	Loose stool, dehydration	Electrolytes, isolate, stop grain	Deworm (Ivermectin, Albendazole), Amprolium (if coccidia), Baycox, Oxytetracycline (if bacterial)	Rehydrate, give B-vitamins, monitor stool
Worms	Pale eyelids, weight loss, rough coat	FAMACHA test, reduce pasture contamination	Ivermectin, Albendazole, Levamisole (rotate), Closantel (for liver fluke)	Repeat in 10–14 days, supplement vitamins
Coccidiosis (Kids)	Diarrhea, weakness, loss of appetite	Electrolytes, isolate	Amprolium 5-day oral, Baycox once	Vitamin B, hygiene, reduce moisture
Bloat	Swollen left side, discomfort	Walk the goat, massage, baking soda drench	Anti-bloat liquid or mineral oil	Feed hay only, avoid sudden diet change
Pneumonia	Coughing, nasal discharge, fever	Shelter, keep warm, isolate	Oxytetracycline or Pen-Strep injections	3–5 days treatment, monitor fever
Mastitis (Does)	Swollen/hard udder, hot to touch	Milk out the infected side, warm compress	Mastitis tubes (e.g. Mastijet), Pen-Strep	Repeat for 3–5 days, hygiene
Foot Rot/Hoof Issues	Limping, swelling, foul smell	Clean and trim hoof	Copper sulfate or formalin footbath, Penicillin	Keep feet dry, clean shed
Orf (Soremouth)	Scabby sores on lips/udders	Isolate, clean sores	Iodine wash, topical antibiotics	Monitor spread, vaccinate if recurring
Pink Eye	Eye tearing, cloudy eye	Isolate, clean eyes	Oxytetracycline spray or injection	Keep in dark if severe, monitor herd

GOAT HEALTH TREATMENT CHECKLIST

Condition	Signs	First Aid	Medication	Follow-Up
Retained Placenta (after kidding)	Placenta not expelled after 24 hrs	Tug gently, give warm water	Oxytetracycline, Uterine boluses	Clean environment, monitor for infection
Ketosis (Pregnancy Toxemia)	Late pregnancy, not eating, weakness	Offer molasses water, propylene glycol	Dextrose IV (vet), Glucose energy boosts	High energy feed, vitamin B
Milk Fever (Calcium Deficiency)	Staggering, tremors, collapse	Warmth, calcium-rich feed	Calcium injection (Calcium Borogluconate)	Gradual diet adjustment
Anaplasmosis/Eperythrozoonosis	Pale eyes, fever, jaundice	Reduce stress, isolate	Oxytetracycline injectable	Monitor herd for ticks/insects
Tick/Flea Infestation	Scratching, hair loss, visible ticks	Spray, dip, isolate	Amitraz dip, Ivermectin injection	Repeat treatment every 14 days
Abscesses (CL or injury)	Lump with pus	Isolate, disinfect, open if mature	Flush with iodine, Penicillin (if infected)	Test for CL if recurring

General Supplements & Support



Item	Use
Vitamin B Complex	Recovery from illness, appetite boost
E-Selen	Muscle support, immunity
Multivitamins	Recovery, stress
Probiotics	After antibiotic use or diarrhea
Molasses + Salt lick	Daily energy and minerals



Farm Tools to Keep Ready

Syringes & needles (3ml, 5ml, 10ml)

Gloves & disinfectants (iodine, Dettol)

Electrolyte mix

Drench gun

Thermometer

Hoof trimmers

Footbath trays

Tick spray / pour-on

Isolation pen

